

BUCKEYE WATCHDOG™ NF



ONE-STEP DISINFECTANT, GERMICIDAL DETERGENT AND DEODORANT



The **WATCHDOG NF** Difference

BUCKEYE WATCHDOG NF is a one-step disinfectant, cleaner and deodorizer designed specifically for the pet care industry. **BUCKEYE WATCHDOG NF** can be used on a variety of hard, nonporous surfaces.

Effective against the following pathogens:

Bordetella bronchiseptica
Salmonella enterica
Staphylococcus aureus

(Animal Viruses:)

- * Avian Infectious Bronchitis virus
- * Avian Influenza A (H5N1) virus
- * Avian Influenza A H3N2 virus
- * Canine Coronavirus
- * Canine Distemper virus
- * Canine Parvovirus – at 4 oz. per gallon
- * Feline Picornavirus
- * Infectious Bovine Rhinotracheitis virus
- * Porcine Parvovirus – at 4 oz. per gallon
- * Pseudorabies virus
- * Transmissible Gastroenteritis virus
- * Vaccinia virus

* Kills Pandemic 2009 H1N1 Influenza A virus.

Available Packaging

W A Q G SS AP 30G 55G 260MP



FEATURES

- Effective against Canine Parvovirus
- pH neutral
- Effective in hard water
- Dilutable at 1 oz. per gallon for normal cleaning, and at 4 oz. per gallon for heavy-duty cleaning and Parvovirus claims

May be used in:

- Veterinary clinics
- Zoos
- Equine farms
- Tack shops
- Pet shops
- Breeding establishments
- Grooming establishments



ONE-STEP DISINFECTANT, GERMICIDAL DETERGENT AND DEODORANT

DIRECTIONS FOR USE:

Preparation of Disinfection/Fungicidal, Virucidal* Use Solution:

1. For heavily soiled areas, a precleaning step is required.
2. Add 1 oz. **BUCKEYE WATCHDOG NF** per gallon of water to disinfect hard, nonporous surfaces. Apply use-solution using a cloth, mop, sponge, sprayer, automatic scrubber, or by immersion so as to thoroughly wet surfaces.
3. Rub with brush, sponge or cloth. Treated surfaces must remain wet for 10 minutes.
4. Rinse or allow to air dry. Rinse all surfaces that come in contact with food with potable water before re-use.

Veterinary Clinics/Breeding Facilities/Zoos/Pet Shops/ Kennels and Grooming Establishments Disinfection Directions:

1. Remove all animals and feeds from areas being treated.
2. Remove all litter and manure from floors, walls and surfaces.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap and detergent, and rinse with water.
5. Apply a use solution of 1 oz. of this product per 1 gallon of water so as to wet thoroughly. Apply by cloth, mop, brush, sponge, automatic scrubber, mechanical spray device or by immersion. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Rub with brush, cloth, or by sponge. Allow surfaces to remain wet for a period of 10 minutes.
6. For Canine Parvovirus saturate all hard, nonporous surfaces with a use-solution of 4 oz. of **BUCKEYE WATCHDOG NF** per gallon of water for a period of 10 minutes.
7. Immerse all halters and other hard, nonporous equipment used in handling and restraining animals as well as forks, shovels and scrapers used in removing litter and manure.
8. Ventilate buildings and other closed spaces.
9. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
10. Thoroughly scrub all treated feed racks, automatic feeders, waterers and other treated equipment that dispenses food or water with soap or detergent, and rinse with potable water before re-use.
11. Change cloth, sponge or towels frequently to avoid redeposition of soil. Prepare a fresh solution at least daily or when use solution becomes visibly dirty.

TECHNICAL SPECIFICATIONS

pH Concentrate	7.6 ± 0.2
pH 1 oz./gal.	7.0 ± 0.2
pH 4 oz./gal.	7.6 ± 0.2
Weight/Gallon	8.34 lbs
Specific Gravity	1.00
Biodegradable	Yes
Concentration of Quat @ 1 oz./gal.	660 ppm

EPA REG. NO. 10324-155-559
EPA EST. NO. 559-MO-1

BUCKEYE INTERNATIONAL, INC.

2700 WAGNER PLACE • MARYLAND HEIGHTS • MO 63043 • USA • 800.321.2583
www.buckeyeinternational.com