

SANICARE TBX™


DISINFECTANT/CLEANER/DEODORIZER



for use against SARS-CoV-2



Available Packaging



The Sanicare TBX Difference

Sanicare TBX is a ready-to-use, quaternary ammonium chloride based product, general non-acid cleaner and disinfectant. It is formulated to disinfect hard, nonporous inanimate environmental surfaces such as floors, countertops, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, bathrooms, shower stalls, bathtubs and cabinets. **For plastic and painted surfaces, spot test on an inconspicuous area before use.** A rinse with potable water is required for surfaces in direct contact with food.

Sanicare TBX also deodorizes those areas that are generally hard to keep smelling fresh, such as garbage storage areas, empty garbage bins and cans, basements, restrooms and other areas which are prone to odors caused by microorganisms.

This product exhibits disinfectant efficacy against Mycobacterium tuberculosis BCG (Mycobacterium bovis) at 20 degrees Centigrade when the treated surface is allowed to remain wet for 5 minutes.

Sanicare TBX is fungicidal against the pathogenic fungus, Trichophyton mentagrophytes (Athlete's Foot Fungus) (cause of ringworm), when used with a **10-minute** contact time on clean, nonporous, hard surfaces as found in bathrooms, shower stalls, locker rooms, or other areas commonly contacted by bare feet. To control mold and mildew on pre-cleaned, hard, nonporous surfaces spray surface to be treated making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth appears.

Sanicare TBX kills the following organisms on hard, nonporous inanimate surfaces:

3 MINUTE KILL TIME FOR BACTERICIDAL ACTIVITY:

Staphylococcus aureus, Salmonella (choleraesuis) enterica, Escherichia coli, Pseudomonas aeruginosa, Escherichia coli 0157:H7, Listeria monocytogenes, Yersinia enterocolitica, Enterococcus faecium, Corynebacterium ammoniagenes, Salmonella (typhi) enterica, and Streptococcus pyogenes.

3 MINUTE KILL TIME FOR ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY:

Methicillin resistant Staphylococcus aureus (MRSA); Vancomycin intermediate resistant Staphylococcus aureus (VISA); Vancomycin resistant Enterococcus faecalis (VRE); Methicillin resistant Staphylococcus epidermidis (MRSE); Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA), (NRS 123) Genotype USA400; Community Associated Methicillin resistant Staphylococcus aureus (CA-MRSA), (NRS 384) Genotype USA300.

KILL TIMES FOR SANICARE TBX when used on environmental, inanimate hard surfaces: Please see attachment. Exhibits virucidal activity against Hepatitis A Virus (HAV), Poliovirus Type 1, Canine Parvovirus, Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Bovine Viral Diarrhea Virus (BVDV), Human Coronavirus, SARS associated Coronavirus, Avian Influenza A strain H3N2, Avian Influenza A strain H9N2, Paramyxovirus (Mumps), Rhinovirus type 39, Rotavirus, HIV-1 (associated with AIDS), Norovirus (Norwalk virus), Feline Calicivirus, and Rabies Virus. See product label for contact times.

SANICARE TBX™

DISINFECTANT/CLEANER/DEODORIZER

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DISINFECTION, DEODORIZING AND CLEANING:

Remove gross filth or heavy soil prior to application of the product. Hold container 6 to 8 inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for **3 minutes** (for Bloodborne Pathogens: allow surfaces to remain wet for HIV-1 **1 minute** and for HBV and HCV **5 minutes**). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum.

One-step cleaning and disinfection of toilet bowls and urinals:

Remove water from bowl by forcing water over trap with a swab mop applicator. Press swab mop against side of the bowl to remove excess water from applicator. Apply 1 to 2 ounces of **Sanicare TBX** evenly, allowing sufficient time for product to cover entire surface of bowl for maximum cleaning. Swab bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Let stand **3 minutes** (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 **1 minute** and for HBV and HCV **5 minutes**) and flush.

EFFICACY TESTS HAVE DEMONSTRATED THAT THIS PRODUCT IS AN EFFECTIVE BACTERICIDE, FUNGICIDE AND VIRUCIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).

TECHNICAL SPECIFICATIONS	
pH	11.7 ± 0.2
Specific Gravity	1.01
Color	Pale Clear Yellow
Odor	Lemon
Freeze-Thaw Stability	Unaffected

EPA REG. NO. 1839-83-559
 DIN: 02258226
 EPA EST. NO. 559-MO-1
 U.S. Patent No. 5,444,094

For more information about Sanicare TBX, scan this code.

