



The **SANICARE D.R.C.** Difference

SANICARE D.R.C. is a ready-to-use, non-acid disinfectant restroom cleaner designed for use on nonporous, hard surfaces. Cleans and deodorizes basins, tubs and glazed tiles. SANICARE D.R.C. is a mildewstat and will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, nonporous surfaces.

Available Packaging



FEATURES

- No measuring or mixing required
- Effectively cleans: Urinals, toilet bowls, sinks, shower stalls, fixtures, floors or any nonporous surface
- Cleans, disinfects and deodorizes in one labor-saving step
- Leaves a clean, fresh scent

SANICARE D.R.C. kills:

(Bacteria:)

- Streptococcus pyogenes
- Salmonella enterica
- Pseudomonas aeruginosa
- Staphylococcus aureus

(Fungi:)

- Trichophyton mentagrophytes (Athlete's Foot Fungus) (Cause of Ringworm)
- Aspergillus niger

(Viruses:)

- *Influenza Type A/Brazil Virus
- *Herpes Simplex Virus Type 2
- *HIV-1 (the AIDS virus)

*Kills Pandemic 2009 H1N1 influenza A virus (formerly called swine flu).



DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DISINFECTION/*VIRUCIDAL/ FUNGICIDAL/ MOLD & MILDEW DIRECTIONS:

If surfaces are visibly dirty preclean first. Apply **SANICARE DISINFECTANT RESTROOM CLEANER** with a mop, sponge, cloth or sprayer on hard, nonporous surfaces to thoroughly wet surfaces. For sprayer applications use a coarse spray device. Spray 6-8 inches from surface until surfaces are thoroughly wet. Avoid breathing spray. Rub with a brush, sponge or cloth. Treated surfaces must remain wet for 10 minutes. Wipe dry with a clean cloth or sponge, or allow to air dry. Rinsing is not necessary on floor surfaces that are to be waxed or polished. For mold and mildew control, repeat application weekly or when growth reappears. Rinse all surfaces that come in contact with food such as countertops, appliances, tables and stovetops with potable water before reuse. This product should not be used to clean utensils, glassware and dishes.

This product is not for use on medical device surfaces.

To Clean and/or Deodorize: Apply by mop, sponge, cloth or spray on hard, nonporous surfaces. Wipe dry with a clean cloth, or sponge, or allow to air dry.

TECHNICAL SPECIFICATIONS	
pН	11.0 ± 0.2
Specific Gravity	1.00
Color	Blue-Green
Fragrance	Floral-Citrus
Freeze/Thaw Stability	Unaffected by freezing
Solvents	None
Free Acids or Alkali	None

EPA REG. NO. 47371-97-559 EPA EST. NO. 559-MO-1 DIN 02040042 **To Clean and Disinfect Toilet Bowls and Urinals:** Remove gross filth from bowl. Remove or expel over the inner trap residual bowl water. Use 4 ounces of product in toilet bowl or urinal. Apply (squeeze) directly around bowl and under rim; then brush or swab thoroughly. Allow product to remain on surface for a contact time of 10 minutes; then flush toilet. This product, when used as directed, will not harm white or colored bowls or inanimate, hard surfaces. Close cap on container after use.

KILLS HIV-1 (AIDS VIRUS) ON PRECLEANED, ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS) OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

Personal Protection: Clean up must always be done wearing protective gloves, gowns, masks, and eye protection.

Cleaning Procedures: Blood and body fluids must be thoroughly cleaned from surfaces and objects before application of disinfectant.

Contact Time: Effective against HIV-1 (AIDS virus) on hard, nonporous surfaces in the presence of a moderate amount of organic soil (5% blood serum) (providing 1800 ppm of active quaternary) for a contact time of 5 minutes at room temperature (20–25° C). Use a ten (10) minute contact time for disinfection against all other bacteria, fungi and viruses claimed (listed on labeling).

Disposal of Infectious Materials: Blood and other body fluids must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.